






nadia30


A standardized and streamlined kit for 3D culture workflows













High-Throughput Solution



High Throughput


Collagen-based scaffolds can be reliably produced on up to 8 Nadia chips in parallel for high-throughput operation

The average scaffold size is ~ 80µm



3D Cell Culture


The collagen-based scaffolds promote cell growth in a 3D physiologically relevant manner.



Single or Multiplets Cell Encapsulation

Dependent on initial cell concentration which is adjustable by the user, more than one cell can be encapsulated in one scaffold

Shown above: co-encapsulation of two differentially-stained cell populations



Cell Release


Viable cells could be released and kept alive for several days after release

Encapsulated cells were cultured and observed over a period of 7 days. Treatment with collagenase led to the digestion of the scaffolds while preserving cell viability.

Kit Benefits



Ideal for 3D culture: the cell physiology is more analogous to that of cells within a living tissue



Compatible with cell release: the nadia3D kit produces collagen-based scaffolds that can be later digested to release cells



No cross-contamination: the nadia3D kit utilizes plastic, single-use cartridges



Standard kit and instructions for making gel scaffolds: no microfluidic knowledge is required



Scalable: choose to run 1, 2, 4 or 8 lanes at once



High throughput
encapsulation of up to 1
million cells in one run

First Steps for 3D Cell Culture



Following sample preparation, collagen-based scaffolds containing cells are produced on the Nadia using the nadia3D kit. The Nadia is an automated, microfluidic droplet-based platform for single cell research that encapsulates up to 8 samples, in parallel, in under 30 minutes and captures up to 1M cells in a single run.

The nadia3D kit creates collagen-based scaffolds, that have distinct advantages over 2D cultures for diverse cell and tissue types as they more closely mimic the natural environments of the cell. As the cell physiology is more analogous to that of cells within a living tissue, the results of assays performed in 3D conditions are more representative of the results you would expect from a living system.

000 «Диаэм»

Москва

ул. Магаданская, д. 7, к. 3 ■ тел./факс: (495) 745-0508 ■ sales@dia-m.ru

www.dia-m.ru

С.-Петербург +7 (812) 372-6040 spb@dia-m.ru

Казань +7(843) 210-2080 kazan@dia-m.ru **Новосибирск** +7(383) 328-0048 nsk@dia-m.ru

Ростов-на-Дону +7 (863) 303-5500 rnd@dia-m.ru **Воронеж** +7 (473) 232-4412

-7 (473) 232-4412 vrn@dia-m.ru

Екатеринбург +7 (912) 658-7606 ekb@dia-m.ru **Йошкар-Ола** +7 (927) 880-3676 nba@dia-m.ru

Kemepobo +7 (923) 158-6753 kemerovo@dia-m.ruu **Красноярск** +7(923) 303-0152 krsk@dia-m.ru

Армения +7 (094) 01-0173 armenia@dia-m.ru

